



# International Journal of Scientific Progress and Research

**ISSN: 2349-4689 (Online)**

Approved By National Science Library (NSL), International Serial Standard and NISCAIR

[Home](#)

[Archives](#)

[Editorial Board](#)

[Review Policy](#)

[Publication Ethics](#)

[Reviewer Guidelines](#)

[Author Guidelines](#)

[Contact Us](#)

**Latest News** IJSPR Volun

[Submit Paper](#)

[Latest News](#)

**Volume 79, October 2021  
Edition Published  
Successfully  
Inviting Research Papers**

IJSPR inviting original research

[IJSPR Released Issues](#)

Issue 188: Vol- 79 No- 10 **NEW**

Issue 187: Vol- 79 No- 09

**Welcome to  
International Journal of Scientific Progress & Research  
ISSN : 2349-4689 (Online)**

**Call For Papers November 2021 (Volume 79) Edition**

**Date of Publication: 30<sup>th</sup> November 2021**

International Journal of Scientific Progress & Research (**IJSPR**) is a peer reviewed monthly online journal. **IJSPR** accepts research works & articles from research scholars, academicians, teachers, professors, doctors, lecturers, and from scientists of their extensive studies and researches. **IJSPR** journal is available for wide range of innovative, intellectual and educational aims to enrich the scientific community all over the world. **IJSPR** releases last week of every month.

The vision and mission of **IJSPR** is to come along with the growing era of engineering tasks, precision, specialized and sensitive devices as well as softwares and simulation tools. are the mandatory use of the short and long term projects. The applications of

[Conferences](#)

- ▶ NCETCE-2018 (Civil) Conference
- ▶ NCETCSE-2018 Conference
- ▶ ETRI-2018 Conference
- ▶ BSIP-2018 Conference
- ▶ NCETCSE-2017 Conference

[Pay Fee](#)

[Impact Factor SJIF 2017](#)

**5.749**

[Download Forms](#)

- ▶ Copyright Form
- ▶ IJSPR Paper Template
- ▶ Membership Form
- ▶ Author Guidelines

## Submit Paper

## IJSR - Issue 131, Volume 45, Number 03

Published On : March 2018

[Latest News](#)
[Previous](#)
[Next](#)

### Volume 79, October 2021 Edition Published Successfully Inviting Research Papers

IJSR inviting original research papers and articles to publish in Volume 79, November 2021 Edition

#### IJSR Released Issues

[Issue 188: Vol- 79 No- 10](#) 
[Issue 187: Vol- 79 No- 09](#)
[Issue 186: Vol- 79 No- 08](#)
[Issue 185: Vol- 79 No- 07](#)
[Issue 184: Vol- 79 No- 06](#)
[Issue 183: Vol- 79 No- 05](#)
[Issue 182: Vol- 79 No- 04](#)
[Issue 181: Vol- 79 No- 03](#)
[Issue 180: Vol- 79 No- 02](#)
[Issue 179: Vol- 79 No- 01](#)
[Issue 178: Vol- 78 No- 01](#)
[Issue 177: Vol- 77 No- 01](#)
[Issue 176: Vol- 76 No- 01](#)
[Issue 175: Vol- 75 No- 01](#)
[Issue 174: Vol- 74 No- 01](#)
[Archive ...](#)

SNo.	Paper Title & Authors	Page
1	<a href="#">Free Trade Agreement and Economic Liberalisation in The Modern Conceptual Ideas Growth and Myth and Reality</a> Sunil. S, Shanimon. S	112-115
2	<a href="#">Ratio Analysis On Working Capital Management of National Hydroelectric Power Corporation Limited Through</a> Hemant Kushwaha	116-120
3	<a href="#">A Study on Consumer Protection Awareness Among Rural Consumer in Karbi Anglong District of Assam</a> Lalit Basumatary	121-124
4	<a href="#">Brain Controlled Robot Using Mind Wave Sensor</a> Mrs. M. Premalatha, P.V. Shamili, P. Sahithi Reddy, S. Sai Krishna, V. Shivateja Reddy	125-129
5	<a href="#">Digital Notice Board Using Raspberry Pi</a> P. Madhusudhan Reddy, M. Himabindhu, R. Ravalireddy, S.Jeevan Kumar	130-134
6	<a href="#">Assessment of Seasonal Variations In Physico-Chemical Properties of Barda Bandharan Wetland : A Limnological Study</a> Dr. Mehul P. Dave, Nikul B. Chavda	135-139
7	<a href="#">Study of A Low Power Pipelined ADC Design</a> Javal Chandravadan Panchal, Prof. Nikhil Ranjan, Prof. K. P. Kanojija	140-143
8	<a href="#">Hydraulic Jack Cum Pipe Bending Machine</a> Abhishek Yadav, Ashish Chourasiya, Deepanshu Shrivastava, Ayush Balagopal, Lakshay Suri, Dheeraj Kumar Raja, Jai Singh Yadav, Prof. Abdul Hameed Khan, Prof. Ashish Kumar Chaturvedi	142-146
9	<a href="#">An Extensive Review on Similarity Validation Based Nonlocal Means Image Denoising</a> Akata Chadar, Versha Sharma	147-152
10	<a href="#">A Grid-Connected New Hybrid Dual Voltage Source Multi-Level Inverter with DC-Fault Blocking Capability and Power Quality Improvement</a> Nithin Nair, Prof. Priyavand Bundela, Prof. Deepika Masand	153-157

[Previous](#)
[Next](#)

#### Download Forms

[Copyright Form](#)
[IJSR Paper Template](#)
[Membership Form](#)
[Author Guidelines](#)

#### Research Fields

[Engineering](#)
[Pharmacy](#)
[Management](#)
[Agriculture Engg.](#)
[Finance/Commerce](#)

**Literature**

**Science**

**Education**

**Engineering Research Streams**

Electronics & Telecommunication

Information Technology

Computer Science Engg.

Mechanical Engg.

Civil Engg.

Chemical Engg.

Electrical Engg.

**Join IJSR**

International Journal of Scientific Progress  
and Research (IJSR) invites Doctors,  
Professors and Researchers to join the  
Editorial and Advisory Board of IJSR.

[Home](#) | [Current Volume](#) | [Research Fields](#) | [Editorial Board](#) | [Free Registration](#) | [E-Certificate](#) | [Contact Us](#)

Copyright © International Journal of Scientific Progress & Research

# Assessment of Seasonal Variations In Physico-Chemical Properties of Barda Bandharan Wetland : A Limnological Study

Dr Mehul P. Dave<sup>1</sup>, Nikul B. Chavda<sup>2</sup>

*Bhakta Kavi Narsinh Mehta University, Junagadh*

*Shri Mahila Arts and Science College, Simar*

**Abstract-Limnology refers to systematic study of plankton and their biodiversity found in aquatic eco-system and their role in maintaining the aqueous ecosystem. Plankton refers to plants and animals that drift with the ocean currents and fresh river water. They are the habitants in the open waters (Limnetic water) of the sea and fresh river water. Phytoplanktons always live near the surface of the sea and fresh water as they require light for photosynthesis, they play important role in transformation of water and carbon dioxide into short chain sugars. The plants in the pelagic zone are exceptionally small, microscopic, and single-celled, buoyantly supported by the density of the surrounding water. Physicochemical parameters are very important factors that play a significant role in wetland plankton diversity and fluctuation in the same. In the present research, we evaluated impact of several abiotics factors on plankton diversity during pre, middle and post winter seasons at of Barada Bandharan wetland, Kodinar, India.**

**Keywords: Plankton Diversity, Abiotic Factor, Limnology, Pelagic zone, Phytoplanktons,**

## I. INTRODUCTION

Planktons are the organisms that live suspended in the water of seas, lakes, ponds, and rivers, and not able to swim against the currents of water. This latter feature distinguishes plankton from nekton, community of actively swimming organisms like fish, larger cephalopods and aquatic mammals. There are two major groups of phytoplankton: (1) non motile, fast-growing diatoms (2) motile flagellates and dinoflagellates, which can migrate vertically in the water column in response to light. Each group exhibits a tremendous variety of cell shapes, many with intricate designs and ornamentations. The diatoms are further divided into two groups based on cell shape: (1) Pennate diatoms, which evolved first during the Late Cretaceous, are long and (2) centric diatoms, which evolved later than the pennates, are shaped like pallboxes and may have elaborate arrays of spines projecting from their cell walls. Phytoplankton has varied in physical and chemical requirements for population growth. Diatoms differ significantly with respect to motility, cell-wall composition and ornamentation, and nutritional and reproductive strategies. Diatoms have cell walls, called

frustules, made of silica (the same material in glass and opal). In contrast, Dino-flagellates can have a rigid cell-wall, called a theca, made of cellulose plates, or they can have a non-rigid cell membrane (no theca). these two forms of Dino-flagellate's structures gave rise to the terms "armored" and "unarmored" (or "naked") Dino-flagellates. Diatoms and Dino-flagellates can be highly ornamented, which aids in species identification. Cell-surface design on some diatoms may help focus light on chloroplast, allowing survival at greater depths where light intensity is very low. Long spines, cell shape, and the formation of chains and colonies make diatoms more difficult for predators to grasp or bite and also assist in flotation. Some Dino-flagellates form chains, whereas others have protuberances that look like wings, crowns, or horns, for similar reasons. Both groups commonly reproduce by simple cell division. Some species of diatoms and Dino-flagellates are known to produce resting stages. Resting spores in diatoms, and cysts in Dino-flagellates, allow species to survive in unfavorable condition. Dino-flagellate's species have feeding veils that are extruded around such food items as diatoms. Both groups are able to absorb nutrients and vitamins into the cell and have distinct preferences for the forms of some of those nutrients.

## II. MATERIALS AND METHODS

### A. Sample collection Points:

Three sample points selected at Barda Bandharan (Sampling Points) with specific GPS location and suitable depth and surface. Sample Collected in plastic bottle (non metallic, free-flushing sample recommended for general purpose of water sampling)

5 liter samples Collected for physicochemical analysis approximately less than 2 feet of river water. Time and temperature measured and transferred all sample as soon as possible to laboratory for study further testing. Temperature range between 18 to 21 °C of samples (Winter Period)

Sr. No.	Barda Bandharan Site Location
1	N 2046 59.4 ,E 070 39 21.1
2	N 2046 59.7 ,E 070 39 27.6
3	N 2046 00.3 ,E 070 39 27.6



Fig.1 : Satellite view of Barda Bandharan Wet land

**B. Bacteriological Analysis:**

Bacteria found in water belong by definition to plankton but because of special techniques required for sampling and identification are considered separately. These organisms are important in the transformation of dead organic materials to inorganic plant nutrients. Some of these marine and freshwater microorganisms (including blue-green algae) fix molecular nitrogen from water containing dissolved air, forming ammonia or related nutrients important for phytoplankton growth. Although little is known about the extent of nitrogen fixation bacteria always are found in water samples. Biological interactions in the ocean are not between populations or between trophic levels, as many box-model representations of pelagic food webs might lead us to think. This allows us to build models and to extrapolate observations beyond the system in which the observations were made. Traditionally, scientists who go on cruises and examine distribution patterns of both biota and environmental properties using sampling are considered biological oceanographers and those who explore the functioning of individuals, for example by conducting laboratory experiments with organisms are considered marine biologists. We need to combine the two approaches to understand the ecology of the oceans.

Standard plate count method used for enumeration of bacteria/biochemical Analysis used for identification of bacterial isolated from water Samples (APHA 2012)

**C. Sample processing for chlorophyll and carotenoids estimation:**

Samples for the chlorophyll estimation were preserved in ice box onboard in darkness to avoid degradation in opaque container covered with aluminum foil. Immediately after reaching the shore ,all samples Put in icebox than transfer all samples as soon as to laboratory for Analysis, 1 liter of collected water sample was filtrate through GF/F filters

(pore size 0.45 um) by using vacuum filtration assembly. After vacuum filtration the glass tissue grinder with glass grinding tube. Glass fiber filter paper will assist breaking the cell during gridding and chlorophyll content was extracted with 10 ml of 90% acetone under cold condition along with saturated magnesium carbonate solution in class screw cap tubes. After an extraction period of 24 hrs, the sample ware transferred to calibrate centrifuge tubes and adjusted the volume to original volume with 90% Acetone .the extract was clarified by using centrifuge in close tubes. The clarified extract was then decanted in clean cuvette and optical density was observed at wavelength according to Chlo. A, B, C and Total chlorophyll with carotenoids. The calculation carryout according to APHA 2012

**D. Sample collection for plankton Analysis**

Collected 1 liter river water sample from three collection site with Plankton net (0.20 microne). After collection of river water samples it's transferred as soon as possible to laboratory for Analysis. Add 4% formalin solution and stay it for 48 hrs, after incubation time period drop count Method used for identified plankton diversity.

**E. Physicochemical Analysis:**

Primary Examination has done Base on Physical examination of water sample by Color, odor and turbidity. pH and Conductivity measured by pH meter and Conductivity meter.

**F. Estimation of Total solid (T.S.)**

Porcelain dish is used for this method; Heat it for 103 to 105 C for 1 hrs. Store and cool dish in desiccators until needed weight immediately before use. (Pre weight) Shake the water sample very well and add 100ml of it in to evaporating Petri dish. Put evaporating dish in to oven at 103 to 105 C for overnight. Next day take out it from oven and cool it in desiccators dish would be having dried residues in it. Measure the weight of evaporating dish. (Post weight)Put the data or pre weight and post weight of the dish in following equation and calculate the amount of total solid present in the sample.

Calculation:  $mg \text{ total solids/L} = (A-B) \cdot 1000 / \text{Sample volume (ml)}$

Where,

A= post weight of dish (weight of dried residues +dish mg)

B= Pre weight (weight of dish mg.)

**G. Estimation of Total dissolved solid (T.D.S.)**

Porcelain dish is used for this method; Heat it for 103 to 105 C for 1 hrs. Store and cool dish in desiccators until

needed weight immediately before use. (Pre weight) Shake the water sample very well and add 100 ml of it in to filtration device that is having glass fiber on it. Apply vacuum and filter out 100ml of sample. Collect the filtrate in to evaporating dish. Put evaporating Petri dish in to oven at 103 to 105 C overnight. Next day take out it from oven and cool it in desiccators dish would be having dried residues in it. Measure the weight of evaporating dish. (Post weight) Put the data of pre weight and post weight of the dish in following equation and calculate the amount of total solid present in the sample.

Calculation:  $\text{mg total dissolved solid/L} = (A-B) \cdot 1000 / \text{sample volume (ml)}$

Where,

A=Post weight of dish

(weight of dried residues+dish, mg)

B=pre weight (weight of dish, mg)

#### H. Estimation of chloride in water sample

Sample preparation: Take 100ml of sample in 250ml conical flask. If chlorine is higher in the sample, dilute the sample and then take 100ml of diluted sample. If the sample is highly colored add 3ml  $\text{Al(OH)}_3$  suspension, mix, settle and filter.

Titration: Set the pH of the sample in the range of 7-10 with the help of  $\text{H}_2\text{SO}_4$  /  $\text{NaOH}$ .

Add 1ml  $\text{K}_2\text{CrO}_4$  indicator solution. Titrate it with standard  $\text{AgNO}_3$  Titrate to a pinkish yellow end point. Be consistent in end point recognition.

Calculation:  $[1] \text{ mg Cl/L} = (A-B) \cdot N \cdot 35450 / \text{ml of sample (100ml)}$

Where, A=ml titration for sample, B=ml titration for blank, C=normality of  $\text{AgNO}_3$  (0.0141N)

$[2] \text{ mg NaCl /L} = (\text{mg Cl/L}) \cdot 1.65$

#### I. Total water hardness:

Take 1ml of water samples than added few drops of the ammonium bisulphate solution add to black-T as indicator. We observed that water sample color is occurrence pink. Then added EDTA slowly drops by drop and water color is blue.

Calculation: Formula:  $1000 \cdot \text{ml of used in EDTA} / \text{ml of water sample}$ .

#### J. Estimation of dissolved oxygen (D.O) and biological oxygen demand (B.O.D)

300 ml of B.O.D. bottle was used for water sample Analysis. In this bottle add 1ml  $\text{MnSO}_4$  solution followed by addition of 1ml alkali iodide acid reagent. Stopper the bottle carefully to exclude and mix by inverting bottle a few times. Add 1ml concentrated  $\text{H}_2\text{SO}_4$ . Res toppe the bottle and mix it thoroughly too completely dissolve the precipitates. Take 200ml of this mixture from bottle to flask.

Add 1ml 2% starch solution as indicator. Titrate it with 0.025  $\text{Na}_2\text{S}_2\text{O}_3$  solutions. Record the end point, when the blue color of starch disappears.

Calculation:  $V_1 \cdot 0.1 \cdot 1000 / 200$  Where,  $v_1$ =Burette no.

#### K. Determination of acidity of water

Hydrogen ions present in a sample as a result of dissociation or hydrolysis of solute react with addition of standard alkali thus acidity depends on end point of the indicator used this colour change of phenolphthalein indicator is used to PH 8.3 at 25°C response to stoichiometric utilization of carbonic acid to bicarbonate.

Mineral acidity:

$$\frac{\text{Volume of NaOH (V1)} * N * 50 * 1000}{\text{Sample taken}}$$

Total acidity

$$\frac{\text{Volume of NaOH (V1)} * N * 50 * 1000}{\text{Sample taken}}$$

### III. RESULTS AND CONCLUSION

Wetland is great ecosystem and its support a great Biodiversity. In present work we investigated interaction between physicochemical parameters with their impact on Microbial biodiversity during pre, middle and post winter time period 2016-2017 of Barda Bandharan wetland (Temporary wetland) Near Barda Village, Kodinar taluka. Microbial biodiversity is the fourth very important factors of ecosystem in water body because its convert complex organic material to simple organic and inorganic compound which utilize by plankton, Water samples collected from wetland of Barda Bandharan around under 2 fit. Included 15 parameters for Analysis like Temperature, pH, Conductivity, T.S, T.D.S., D.O., B.O.D. water Hardness and chloride. Temperature range 19.0°C to 20.5°C. pH range of wetland water was 7.96 to 8.9, pH of water samples were normal Range as per standard but higher pH of water noted on

Nove-11 Month. Higher results of pH indicated salts concentration may higher. Conductivity of water were higher in the sample (Dec-21), higher conductivity indicates salts concentration higher in water sample. Dissolved oxygen (D.O) and Biological oxygen demand (B.O.D) data indicated that dissolve oxygen level range 7.9 to 8.9 in water. Higher D.O. value indicates good condition for aquatic life inside the water. T.S. and T.D.S. data of water samples were higher and fluctuate more during time period. T.S.range of sample 1000 mg/L to 1920 mg/L, Higher, TDS of samples range 300 mg/L to 490 mg/L the data of T.S and T.D.S is higher than normal range its indicated water is not directly use for Agriculture and drinking purpose, higher values is also dangerous for normal aquatic life. Water hardness is another parameter which indicated salts quality in water samples like carbonate and many other salts in water sample. Water hardness Ranges were 300 mg/L to 670 mg/lit, salt concentration were increase during sampling time period (Table:01 and Figure:01 to 08)

we investigated hydrology parameter influence the biodiversity and many other biological parameters, microorganisms also play in maintain organic and inorganic compound in water ecosystem, from above research case study we conclude that plankton and microbial interaction play important role in maintain wetland ecosystem.

Biological Parameter included chlorophyll estimation in which Analysis of Chlo A, chlo B and Total Chlorophyll with cartenoids concentration. Chlorophyll play important role in production of organic molecules in water body ecosystem and it's maintaining food web chain in water body ecosystem. Plankton Analysis in which Zooplankton and phytoplankton are very important biotic factor maintain water body ecosystem. Phytoplankton is primary food producer which is consuming by zooplankton and fish with many other water body animals depend on zooplankton concentration in wetland ecosystem.

Chlorophyll estimation of collected wetland water was indicating that total chlo. Concentration increased during data Analysis with Time period, its indicated organic concentration increase during time period and due to that Bacterial concentration may increase.

Concentration of chlorophyll also indicated increase concentration of phytoplankton, phytoplankton are primary producer of food in water ecosystem they play a major role in maintain water ecosystem. We isolated 36 spp. of phytoplankton during Analysis; Ditoms concentration was higher in which *Rhizosolenia setigera* was predominant in water system. *Cylindrotheca* spp. was also found during water sample analysis and cynobacteria were predominant

found in wetland water system they also important role in photosynthesis.

Plankton Analysis in which Zooplankton and phytoplankton are very important biotic factor maintain water body ecosystem. Phytoplankton is primary food producer which is consuming by zooplankton and fish with many other water body animals depend on zooplankton concentration in wetland ecosystem. Concentration of chlorophyll also indicated increase concentration of phytoplankton, phytoplankton are primary producer of food in water ecosystem they play a major role in maintain water ecosystem. Isolated 18 spp. of Zooplankton during Analysis; *Acrocalanus longicornis* concentration was higher in which copepod nauplius was predominant in water system. *Cylindrotheca* spp. was also found during water sample analysis and cynobacteria were predominant found in wetland water system they also important role in photosynthesis.

## REFERENCES

- [1] American Public Health Association.2012. Standard methods for the examination of water and wastewater.18th edition. American Public Health Association. Washington, D.C., USA.
- [2] Attayde, J. L., and R. L. Bozelli. 1998. Assessing the indicator properties of zooplankton assemblages to disturbance gradients by canonical correspondence analysis. *Canadian Journal of Fisheries and Aquatic Sciences* **55**:1789–1797.
- [3] Balcer, M. D., N. L. Korda, and S. I. Dodson. 1984. Zooplankton of the Great Lakes. University of Wisconsin Press, Madison, Wisconsin, USA.
- [4] Bertram, P., and N. Statler-Salt. 1999. Selection of indicators for Great Lakes basin ecosystem health. Environment Canada and U.S. Environmental Protection Agency. State of the Lakes Ecosystem Conference 1998. [Online, URL: [www.cciw.ca/solec&.](http://www.cciw.ca/solec&.)]
- [5] Berzins, B., and J. Bertilsson. 1989. On limnic microcrustaceans and trophic degree. *Hydrobiologia* **185**:95–100.
- [6] Berzins, B., and B. Pejler. 1989. Rotifer occurrence and trophic degree. *Hydrobiologia* **182**:171–180.
- [7] Botts, P. S. 1999. Lake Erie coastal wetlands: a review and case study of Presque Isle invertebrates. Pages 995–1012 in
- [8] E. Keas, and C. A. Stricker. 1999. Development of a preliminary invertebrate index of biotic integrity for Lake Huron coastal wetlands. *Wetlands* **19**:869–882.
- [9] Campbell, J. M. 1993. The cladoceran species of inshore habitats of Lake Erie at Presque Isle. *Journal of the Pennsylvania Academy of Science* **67**:115–119.
- [10] Cardinale, B. J., V. J. Brady, and T. M. Burton. 1998. Changes in the abundance and diversity of coastal wetland fauna from the open water/macrophyte edge towards shore. *Wetlands Ecology and Management* **6**:59–68.
- [11] Chow-Fraser, P. 1998. A conceptual model to aid restoration of Cootes Paradise Marsh, a degraded coastal wetland of Lake Ontario, Canada. *Wetland Ecology and Management* **6**:43–57.

- [12] Chow-Fraser, P. 1999. Seasonal, interannual and spatial variability in the concentrations of total suspended solids in a degraded coastal wetland of Lake Ontario. *Journal of Great Lakes Research* **25**:799–813.
- [13] Chow-Fraser, P., and D. A. Albert. 1999. Coastal wetland ecosystems: biodiversity investment areas. EnvironmentCanada and U.S. Environmental Protection Agency. State of the Lakes Ecosystem Conference 1998.[Online, URL:[^www.cciw.ca/solec&.](http://www.cciw.ca/solec&.)].
- [14] Chow-Fraser, P., V. Lougheed, V. Le Thiec, B. Crosbie, L.Simser, and J. Lord. 1998. Long-term response of the biotic community to fluctuating water levels and changes in water quality in Cootes Paradise Marsh, a degraded coastal wetland of Lake Ontario. *Wetland Ecology and Management* **6**:19–42.